
Extreme Quasar Variability

Daniel Stern
(Jet Propulsion Laboratory/
California Institute of Technology)




Matthew J. Graham (California Institute of Technology)




- quasars can be distinguished from Galactic stars based on their optical light curves
- quasar light curves well-described by a "damped random walk"


PG 1302-105




- Extreme Activity I: periodic quasars
- binary supermassive black holes???




only ~1/10,000 quasars should show this much optical variability based on DRW model



- Extreme Activity 2: changing look/changing state quasars (step function lightcurve)
- e.g., FeLoBAL turning into a normal BAL quasar over a decade




- Extreme Activity 2: changing look/changing state quasars (step function lightcurve)
- e.g., WISE-selected changing state quasars





• Extreme Activity 3: Tidal Disruption Events in AGN


IES 1927+65



GraL Collaboration




Alberto Krone-Martins (UC-Irvine)






S.G. Djorgovski (Caltech)

• Extreme Activity 4: Strong Gravitational Lenses






- Extreme Activity 5: AGN flares
- obviously diverse flare morphologies = diverse physics


THE ASTROPHYSICAL JOURNAL LETTERS


Ram-pressure Stripping of a Kicked Hill Sphere: Prompt Electromagnetic Emission from the Merger of Stellar Mass Black Holes in an AGN Accretion Disk

```
B. McKernan<sup>1,2,3</sup> D, K. E. S. Ford<sup>1,2,3</sup>, I. Bartos<sup>4</sup> D, M. J. Graham<sup>5</sup> D, W. Lyra<sup>6,7</sup> D, S. Marka<sup>8</sup>, Z. Marka<sup>8</sup>, N. P. Ross<sup>9</sup>, D. Stern<sup>7</sup> D, and Y. Yang<sup>4</sup> Published 2019 October 17 • © 2019. The American Astronomical Society. All rights reserved. The Astrophysical Journal Letters, Volume 884, Number 2
```



Masses in the Stellar Graveyard






CASTOR: the Cosmological Advanced Survey Telescope for Optical and ultraviolet Research








- I-meter unobscured aperture
- wide field-of-view $(0.61^{\circ} \times 0.38^{\circ}; FWHM = 0.15'')$
- 3-band imaging + spectroscopy
- planned launch in ~2027
- broad science, from exoplanets to cosmology
- 8 science working groups, including both Time Domain Astrophysics and AGN
- planned wide-field blind time-domain legacy survey, as well as targeted time-domain follow-up of AGN
 https://www.castormission.org/

UVEX: Ultraviolet Explorer



- Pl: Fiona Harrison (Caltech)
- Project Scientist: Brian Grefenstette (Caltech)
- will do a multiple-cadence, all-sky UV survey, reaching 50-100x deeper than GALEX
- 75 cm effective aperture
- wide field-of-view $(3.5^{\circ} \times 3.5^{\circ})$
- FWHM ≤ 2.25"
- 2-band imaging (NUV + FUV) + longslit spectroscopy
- if selected, planned launch in ~2028







THEEND



BACK-UP SLIDES



A Black Hole Over the Limit





Whereas Chris Belczynski and Daniel Holz believe that astrophysical black holes should not exist in the mass range between 55 and 130 solar masses because of pair instability; and whereas Carl Rodriguez and Sourav Chatterjee believe that such black holy would form in dynamical environments and continue to participate in mergers;

they wager, a \$100 bottle of wine that, within the first 100 Gw... compact binary coalescence detections, at least one will have a component in the 55-130 Morange. If individual events have mass ranges straddling the interval boundary, the betting parties agree that Ilya interval will serve as an arbiter of the statistical evidence. Mandel will serve as an arbiter of the statistical evidence.

chis belighti,
1/Chris Belczynski/
1/Oaniel Holt/

Signed in Aspen, Co, on 10 Feb. 2017

Carl Rodriguery

/Carl Rodriguer/

Somar Chatterine

/Sourau Chatterine

/Witnessed by Ilya Mandel/ Hettaro

Tel (970) 925-2585 • Fax (970) 920-1167 • ACP@aspenphys.org • www.aspenphys.org

Potential false positives:

AGN variability: Analyzed light curves of 2.5 million WISE-selected AGN from Assef et al. (2018) with ZTF coverage, looking for flares. Find 393 events. Comparison with damped random walk (DRW) model to ZTF light curves and identifying sources where flare is strong preferred over regular quasar variability drops this number to 13. So probability of a linear light curve with a well-detected >20 day flare is ~5e-6.

Look elsewhere effect: Looked at longer term CRTS light curves for all 3255 AGN within GW 190521g area, and Monte Carlo'd their light curves (1000x per AGN), assuming DRW model, with ZTF sampling. Find just 5 comparable flares, without visual inspection. So O(0.5%) likelihood of chance event.

Supernova: Rejected given the timescale (too short for a SN) and uniform color over time (SNe vary in color as they evolve).

Microlensing: Rejected given the level of magnification, expected years-long event, not weeks.

Tidal disruption event (TDE): Rejected; doesn't match luminosities and GW signal.

Predictions:

EMGW from binary BH mergers: Preferentially should be detected for more massive BHs. Would be good if O4+ include (crude) BH masses.

Off-center flare: If got spectra soon enough, expect signatures from non-uniform illumination of the BLR.

Return to the disk: Expect BH to return to the disk, assuming typical kicks. So more BHL emission in ~1.5 years.

Working on longer, more detailed paper(s) including full GW event sample and more details on the predicted light curves.

Assuming a cosmology, can get constraints on orientation of the BBH merger (= AGN orientation). Or, if you're more bold, assuming AGN orientation, get cosmology.

Extreme Quasar Variability

Daniel Stern
(Jet Propulsion Laboratory/
California Institute of Technology)



Matthew J. Graham (California Institute of Technology)

Extreme Quasar Variability

Daniel Stern
(Jet Propulsion Laboratory/
California Institute of Technology)



Matthew J. Graham (California Institute of Technology)